

A Deeper Look into Indianapolis: CO₂ **Emissions and Economic Conditions**



Joan Haling, Kenneth Davis, Thomas Lauvaux, Zach Barkley The Pennsylvania State University July 30th, 2015

Abstract

Major cities across the United States account for nearly one-quarter of all global carbon dioxide emissions. Therefore Arizona State University researchers have set out to create a system that allows for all fossil fuel carbon dioxide emissions to be quantified at the scale of buildings, streets and factories. The system, named "Hestia", was first tested in Indianapolis, Indiana. In an attempt to better understand this first case study, the economic conditions of Indianapolis were researched and, in an exploration of the data product, the results from Hestia were considered and analyzed to better understand their contents. This showed Indianapolis to have a growing economy and emissions that change based on a

Introduction

The Hestia Project builds on the Vulcan Project, which quantified fossil fuel CO2 in the United States at sub-county spatial resolution and temporal resolution for the year 2002 (Gurney, et al. 2009). Hestia has taken that information and adapted it into a bottom-up approach that models fossil fuel carbon dioxide emissions at the scale of individual buildings. road segments as well as industrial and electricity production facilities (Gurney et al, 2012). The data collected and analyzed from the Hestia Project encompasses the years 2002 to 2013 and ranges over eight different sectors: airport, commercial, industrial, mobility, non-road, residential, utility and rail.



Focus Questions


Focus questions:

- I. How have the economic conditions in Indianapolis changed overtime and how do they compare with other major United States cities?
- II. What do the carbon dioxide emission results from the Hestia Project show?
- III. What can the results from the Hestia Project help us understand about emission changes by region and sector?


- Methods
 The economic information came from StatsIndiana and the U.S. Department of Commerce and then was loaded on the Matlab to find comparisons and
- The heating degree days were calculated by information presented on Weather Underground.


Using the given Hestia data the carbon dioxide emission results were analyzed on Matlab to find ratio changes between the different years.

Results




Fopulation, real GDP and CU2 emissions as a fraction of the mean Population and real GDP show continuous data while CO₁ emissions are from year 2002, 2010, one million- according to 2010 census





III. Emissions by Region and Sector



Discussion and Conclusions

Economic Conditions


- Indianapolis MSA has an economy that is slightly above average with an overall growing population, GDP and rate of CO2 emissions.
- The growth of rate CO₂ emissions is correlated with the growth rate of GDP. (GDP used to compute CO₂?)

II. Emissions Results

- · The Hestia Project is a modelling program that takes the carbon dioxide emission data from 2002 and adjusts the yearly results in accordance to factors such as GDP.
- The total amount of carbon dioxide emissions in 2012 was lower because there was a warm winter and therefore the residential sector did not need to consume as much heat.

III. Emissions by Region and Sector

- The 2013/2002 ratio carbon dioxide emissions plot showed county level changes, with the most detail being in Marion County and the largest increase in Boone County.
- The NonRoad sector plot showed an increase in the level of emissions in Boone County. Causes for this increase over the 11 year period could be an increase in construction or a problem with the model.
- Results show that while there was a dip in emissions in 2012 due to the Residential sector, overall emissions increased during the 11 years which is largely due to the OnRoad (mobility) sector.



References

story for Indianapolis, IN | Weather Underground." Weather Underground. The Weather Channel, 2015. Web. 2 July 2015.

Acknowledgements

This work was [partially] supported by the National Science Foundation through the Network for Sustainable Climate Risk Management (SCRiM) under NSF cooperative agreement GEO-1240507. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.